arXiv:0810.2597v3 [math.AG] 30 Aug 2009

STRONG RATIONAL CONNECTEDNESS OF SURFACES

CHENYANG XU

Abstract. This paper focuses on the study of the strong rational connectedness of smooth rationally connected surfaces. In particular, we show that the smooth locus of a log del Pezzo surface is strongly rationally connected. This confirms a conjecture due to Hassett and Tschinkel in [HT08].

Contents

1. Introduction 1
2. Twisted curves and twisted stable maps 4
3. Smoothing 8
4. C-noncontracted minimal model program 13
5. Nonproperness of the complement of the very free locus 17
6. Examples 20

References 23

1. Introduction

Throughout this paper, except in Section 2 and 6, the ground field k will be an algebraically closed field of characteristic zero.

1.1. **Definition.** A variety X is called *rationally connected* if there is a family of proper rational curves $g : U \to Z$ and a cycle morphism $u : U \to X$ such that $u^{(2)} : U \times_Z U \to X \times_k X$ is dominant.

We refer to [Ko96] for the background of rationally connected varieties. Given a smooth variety X, it is rationally connected if and only if there is a morphism $f : \mathbb{P}^1 \to X$ such that f^*T_X is ample (cf. [Ko96] IV.3.3.1 and 3.7). Furthermore, it is known (cf. ibid IV.3.9.4) that for any smooth rationally connected variety X over k, there is a unique largest nonempty open subset $X^{vf} \subset X$ with the property that $x \in X^{vf}$ if and only if there is a morphism $f : \mathbb{P}^1 \to X^{vf}$ such that:

(1) $x \in f(\mathbb{P}^1)$;
(2) $f^*(T_X)$ is ample.

We call X^{vf} the *very free locus* of X. If $X^{vf} = X$, X is called *strongly rationally connected* (cf. [HT08], 14).

1.2. **Theorem** ([KMM92b] 2.1, [Ko96] IV.3.9). *A proper and smooth rationally connected variety X/k is strongly rationally connected.*

A result of Campana and Kollár-Miyaoka-Mori states that all smooth Fano varieties are rationally connected ([Ca92], [KMM92a] or [Ko96]). More generally, log $\mathbb{Q}$-Fano varieties are rationally connected ([Zh06] or [HM07]). A *log $\mathbb{Q}$-Fano variety* (X, Δ) is a normal projective variety X with a Weil divisor Δ such that (X, Δ) is *klt* and $K_X + \Delta$ is anti-ample. A log $\mathbb{Q}$-Fano surface is also called a *log del Pezzo surface.*

Given a log $\mathbb{Q}$-Fano variety (X, Δ), one can ask whether the smooth locus X^{sm} of the underlying space X is rationally connected. In general determining the rational connectedness of a nonproper variety can be formidable, and the answer to this question is not known except for the surface case.

1.3. **Theorem** ([KM99], 1.6). *If (S, Δ) is a log del Pezzo surface, then the smooth locus S^{sm} of S is rationally connected.*

If we drop the properness assumption, it is not known whether the rational connectedness implies the strong rational connectedness ([Ko96], IV.3.9.5). The main purpose of this paper is to show that the smooth loci of log del Pezzo surfaces are not only rationally connected but also strongly rationally connected. This confirms a conjecture due to Hassett and Tschinkel ([HT08], Conjecture 19).

1.4. **Theorem.** *(Notation as above.) S^{sm} is strongly rationally connected.*

Applying the same technique, we can prove

1.5. **Theorem.** *Let S be a projective surface which has at worst Du Val singularities. If S^{sm} is rationally connected, then S^{sm} is indeed strongly rationally connected.*

The main strategy to prove (1.4) consists of two steps: first, using an argument from the Minimal Model Program(MMP), and applying Keel-M^{c}kernan's result (1.3), we prove (4.4) saying that there are at most finitely many points in S^{sm} which are not in the very free locus. In the second step, we need to prove that indeed S^{sm} is equal to its very free locus. Given two smooth points p and q of S, we choose two sequences of general points to specialize to them. Applying Keel-M^{c}Kernan's theorem again, we can connect any two general points by a proper rational curve contained in S^{sm}. Then we want to construct a limiting object containing p and q, and show that after adding very free curves to the limiting object, we can smooth it away from the singular points of S, but still containing p and q. For the purpose of doing deformation, we will present two approaches. The first approach is studying rational curves on the smooth stack Deligne-Mumford $\mathcal{S}$, which realizes S as its coarse moduli space. The theory of the moduli space of twisted stable maps which have Deligne-Mumford stacks as the target spaces (cf.[AV02]) provides the framework to study this. The advantage of considering $\mathcal{S}$ is that its smoothness allows us to do the deformation theory in a similar way as the classical case when the target space is a smooth variety (See Section 2 and 3). We get the limiting object, which could be reducible and twisted, from the properness of $\mathcal{K}_{0,2}(\mathcal{S}, d)$. The first step allows us to attach teeth to general points in each component of the image of the limiting object. By imitating the analogue argument for the classical curves, we show that after attaching enough teeth, it eliminates all obstructions of deforming the morphism with a finite set of points fixed. We thereby can smooth the limiting object, with the additional property that the image still contains the two given points. Then the smoothing curve is a very free curve, which we can choose to be contained entirely in the scheme locus of $\mathcal{S}$ (See (3.8)).

The second approach is similar. But instead of considering the stack covering, it uses the new concept of *pseudo-ideal sheaves* introduced in [SR09]. The argument is due to Jason Starr, who was inspired by reading the first version of this note. In fact, the following stronger result is established by using pseudo-ideal sheaves,

1.6. **Theorem.** *(Starr) If X^0/k is a smooth rationally connected variety (of arbitrary dimension), $W \subset X^0$ is a closed subscheme which is proper over k satisfying that $U := X^0 \setminus W$ is strongly rationally connected, then X^0 itself is strongly rationally connected.*

For some special log del Pezzo surfaces, the strong rational connectedness of the smooth loci has been established in [HT08] and [Kn08]. Then these results have been applied to prove certain weak approximation results for rational surfaces over the function field of curves. In [Xu], we will use the results of this note to establish weak approximations for more del Pezzo surfaces.

From our argument, it is also interesting to ask the following question: if we assume U to be a nonproper smooth rationally connected surface, and $S \supset U$ a normal compactification, then what type of singularities can S have? We give the following answer:

1.7. **Theorem.** *(Notation as above). S contains only rational singularities.*

The content of this note is organized as follows: in Sections 2 and 3, we develop a twisted version of the classical theory of morphisms from curves to varieties; then we apply this tool to prove our main theorems (1.4) and (1.5) in Section 4; in Section 5, we very briefly sketch the theory of pseudo-idea sheaves and then present the proof of (1.6); in section 6, we give two examples: the first one shows that MMP does not always preserve the rational connectedness of the smooth locus; the second one is due to Kollár which yields a del Pezzo surface with A_1 singularities in characteristic 2, whose smooth locus does not contain any free curve.

Acknowledgement: I would like to thank Igor Dolgachev, Brendan Hassett, Amit Hogadi, Amanda Knecht, János Kollár for useful conversations and emails. I especially want to thank Dan Abramovich for suggesting me a crucial idea to prove Theorem (1.4) and Jason Starr for allowing me to use his unpublished arguments. Thanks also to Luli Jiarong for his help on English; any remaining mistakes are my own. The author is partially supported by Clay liftoff fellowship. Part of the work was done when the author visited Universität Duisburg-Essen. The author wishes to thank Hélène Esnault for her hospitality during the visit. This material is also based upon work when the author is in Institute for Advanced Study and supported by the NSF under agreement No. DMS-0635607.

2. Twisted curves and twisted stable maps

In this section, we give a short introduction to the theory of twisted curves and n-pointed twisted stable maps. We refer to [AV02] and [Ol07] for more details. Let S be a scheme and $\mathcal{C}/S$ a proper flat tame Deligne-Mumford stack $\mathcal{C} \to S$ whose fibers are purely one-dimensional and geometrically connected with at most nodal singularities (recall that $\mathcal{C}$ is tame if for every algebraically closed field k and morphism $x : \mathrm{Spec}\,(k) \to \mathcal{C}$ the stabilizer group $\mathrm{Stab}_{\mathcal{C}}(x)(k)$ has order invertible in k). Let $\mathcal{C} \to C$ be the coarse moduli space of $\mathcal{C}$, and let $C^{sm} \subset C$ be the open subset where $C \to S$ is smooth. Assume that the inverse image $\mathcal{C} \times_C C^{sm} \subset \mathcal{C}$ is equal to the open substack of $\mathcal{C}$ where $\mathcal{C} \to S$ is smooth and that for every geometric point $\bar{s} \to S$ the map $\mathcal{C}_{\bar{s}} \to C_s$ is an isomorphism over some dense open subset of $\mathcal{C}_{\bar{s}}$. Then the coarse space C is a nodal curve over S, and it is well-known that, for any geometric point mapping to a node $\bar{s} \to C$ there exists an étale neighborhood $\mathrm{Spec}\,(A) \to C$ of $\bar{s}$ and an étale morphism

$$\mathrm{Spec}\,(A) \to \mathrm{Spec}\,_S(\mathcal{O}_S[x, y]/xy - t)$$

for some $t \in \mathcal{O}_S$, such that the pullback $\mathcal{C} \times_C \mathrm{Spec}\,(A)$ is isomorphic to

$$[\mathrm{Spec}\,(A[z, w]/zw = t', z^n = x, w^n = y)/\Gamma]$$

for some element $t' \in \mathcal{O}_S$, where Γ is a finite cyclic group of order n invertible in A such that if $\gamma \in \Gamma$ is a generator then $\gamma(z) = \xi z$ and $\gamma(w) = \xi' w$ for some primitive *nth* roots of unity ξ and ξ'. The stack is called *balanced* if étale locally there exists such a description with $\xi' = \xi^{-1}$.

2.1. **Definition** (twisted curve)**.** ([AV02], 4.1.2). A *twisted curve* is a stack $\mathcal{C} \to S$ as above such that the action at each nodal point $\bar{s} \to \mathcal{C}$ is balanced. A twisted curve $\mathcal{C} \to S$ has genus g if the genus of $C_{\bar{s}}$ is g for every geometric point $\bar{s} \to S$. An *n-pointed twisted curve* is a twisted curve $\mathcal{C} \to S$ together with a collection of disjoint closed substacks $\{\Sigma_i\}_{i=1}^n$ of $\mathcal{C}$ such that:

(1) each $\Sigma_i \subset \mathcal{C}$ is contained in the smooth locus of $\mathcal{C} \to S$;
(2) the stacks Σ_i are étale gerbes over S;
(3) if $\mathcal{C}_{gen}$ denotes the complement of the Σ_i in the smooth locus of $\mathcal{C} \to S$, then $\mathcal{C}_{gen}$ is a scheme.

2.2. **Remark.** Unlike [AV02], we will only use balanced twisted curves in this paper, so we usually omit the adjective "balanced". Abramovich and Vistoli prove that the moduli stack parametrizing the twisted stable maps from balanced twisted curves is an open and closed substack of the stack parametrizing all twisted stable maps (cf. [AV02] 8.1.1).

In [Ol07], Olsson got the following description of the versal deformation of a given n-pointed twisted curve. Let $\mathcal{M}^{tw}_{g,n}$ denote the fibered category over $\mathbb{Z}$ which to any scheme T associates the groupoid of n-marked genus g twisted curves $(\mathcal{C}, \Sigma_i)$ over T, and let $\mathcal{S}_{g,n}$ denote the fibered category over $\mathbb{Z}$ which to any scheme S associates the groupoid of all (not necessarily stable) n-pointed genus g nodal curves $\mathcal{C}/S$. Now consider a field k and an object $(\mathcal{C}, \Sigma_i) \in \mathcal{M}^{tw}_{g,n}(k)$. Let (C, σ_i) be the coarse moduli space, and let R be a versal deformation space for the object $(C, \sigma_i) \in \mathcal{S}_{g,n}(k)$. Let $q_1, ..., q_m \in C$ be the nodes, r_i the order of the stabilizer group of a point of C lying above q_i. It is well-known (cf. [DM69]) that there is a smooth divisor $D_i \subset \mathrm{Spec}\,(R)$ classifying deformations, where q_i remains a node. In other words, if $t_i \in R$ is an element defining D_i then in an étale neighborhood of q_i the versal deformation $\widetilde{C} \to \mathrm{Spec}\,(R)$ of (C, σ_i) is isomorphic to

$$\mathrm{Spec}\,(R[x, y]/xy - t_i).$$

As a corollary to the above result, Olsson shows

2.3. **Proposition** ([Ol07], 1.11). *(Notations as above.) A versal deformation space for the twisted curve* $(\mathcal{C}, \Sigma_i)$ *is given by*

$$R[z_1, ..., z_m]/(z_1^{r_1} - t_1, ..., z_m^{r_m} - t_m).$$

In our later discussion, we will be interested in the smooth objects of the versal family. From this proposition, we know that such objects always exist.

Now we consider a proper tame Deligne-Mumford stack $\mathcal{M}$ admitting a projective coarse moduli scheme M. We fix an ample invertible sheaf on M. To compactify the n-pointed stable maps from curves to a Deligne-Mumford stack, Abramovich and Vistoli made the important observation that we have to enlarge the source from the category of curves to the category of twisted curves. (The collection of twisted curves over a scheme S form a 2-category which is in fact equivalent to a 1-category (cf. [AV02], 4.4.2).)

2.4. **Definition** (n-pointed twisted stable map). A twisted stable n-pointed map of genus g and degree d over S $(\mathcal{C} \to S, \Sigma_i^{\mathcal{C}} \subset \mathcal{C}, f : \mathcal{C} \to M)$ consists of a commutative diagram

$$\begin{tikzcd}
\mathcal{C} \arrow[r] \arrow[d] & \mathcal{M} \arrow[d] \\
C \arrow[r] \arrow[d] & M \\
S &
\end{tikzcd}$$

along with n closed substacks $\Sigma_i^{\mathcal{C}} \subset \mathcal{C}$, satisfying:

(1) $\mathcal{C} \to C \to S$ along with $\Sigma_i^{\mathcal{C}}$ is a twisted nodal n-pointed curve over S,
(2) the morphism $\mathcal{C} \to \mathcal{M}$ is representable, and
(3) $(C \to S, \Sigma_i^C, f : C \to M)$ is a stable n-pointed map of degree d.

2.5. **Theorem** ([AV02], 1.4.1)**.** *Let* $\mathcal{K}_{g,n}(\mathcal{M}, d)$ *be fibered over* Sch/S*, the category of the twisted stable n-pointed maps* $\mathcal{C} \to \mathcal{M}$ *of genus* g *and degree* d*.*

(1) *The category* $\mathcal{K}_{g,n}(\mathcal{M}, d)$ *is a proper algebraic stack.*
(2) *The coarse moduli space* $K_{g,n}(\mathcal{M}, d)$ *of* $\mathcal{K}_{g,n}(\mathcal{M}, d)$ *is projective.*
(3) *There is a commutative diagram*

$$\begin{tikzcd}
\mathcal{K}_{g,n}(\mathcal{M}, d) \arrow[r] \arrow[d] & \mathcal{K}_{g,n}(M, d) \arrow[d] \\
K_{g,n}(\mathcal{M}, d) \arrow[r] & K_{g,n}(\mathcal{M}, d),
\end{tikzcd}$$

where the top arrow is proper, quasifinite, relatively of Deligne-Mumford type and tame, and the bottom arrow is finite. In particular, if $\mathcal{K}_{g,n}(M, d)$ *is a Deligne-Mumford stack, so is* $\mathcal{K}_{g,n}(\mathcal{M}, d)$*.*

In our paper, we also need to study morphisms between stacks and their deformation theory, which has been intensively studied in [Ol06a], [Ol06b]. From Olsson's results, we can see that when the stacks are Deligne-Mumford, the theory is similar to the case of schemes. For the reader's convenience, we summarize the results which we need in our proof,

2.6. **Theorem.** *Let* $\mathcal{C}/S$ *be a twisted curve and* $\mathcal{Y}/S$ *a Deligne-Mumford stack.* $\underline{\mathrm{Hom}}_S(\mathcal{C}, \mathcal{Y})$ *is represented by a Deligne-Mumford stack. The fibered subcategory of representable morphisms*

$$\underline{\mathrm{Hom}}_S^{rep}(\mathcal{C}, \mathcal{Y}) \subset \underline{\mathrm{Hom}}_S(\mathcal{C}, \mathcal{Y})$$

is an open substack.

Proof. See the proof of ([Ol07], 1.15). □

2.7. **Theorem.** *Let* k *be a field. In the above notations, choose* $S = \mathrm{Spec}\,(A)$*, where* A *is an artinian ring with residue field* k*. Assume* $g : \mathcal{C}/A \to \mathcal{Y}/A$ *a representable morphism. Denote* $L_{\mathcal{C}/\mathcal{Y}}$ *as the contangent complex of* g*. If* $\gamma :$ $\mathrm{Spec}\,(A) \to \mathrm{Spec}\,(A')$ *is a closed immersion defined by a square-zero idea* $I \subset A'$*, and* $i : \mathcal{C}/A \to \mathcal{C}'/A'$ *is a flat extension with an isomorphism* $\mathcal{C}' \times_{A'} A \cong \mathcal{C}$*, then*

(1) *There is a canonical obstruction class* $o(g, i) \in \mathrm{Ext}^1(Lg^*(L_{\mathcal{Y}/A}), I)$ *whose vanishing is necessary and sufficient for the existence of an arrow* $g' :$ $\mathcal{C}' \to \mathcal{Y}$ *extending* g*.*
(2) *If* $o(g, i) = 0$*, then the set of isomorphism classes of maps* $g' : \mathcal{C}' \to \mathcal{Y}$ *as in (1) is naturally a torsor under* $\mathrm{Ext}^0(Lg^*(L_{\mathcal{Y}/A}), I)$*.*

Proof. See the proof of [Ol06a], Theorem 1.5 (i) and (ii)]. □

2.8. **Remark.** Like ([Ol06a], 1.7), Theorem (2.7) can be used to define an obstruction theory in the sense of ([Ar74], 2.6). Recall that such a theory consists of the following (in the following all rings are over S):

(1) For every surjection of noetherian rings $A \to A_0$ with nilpotent kernel and $a \in \underline{\mathrm{Hom}}_S^{rep}(\mathcal{C}, \mathcal{Y})(A)$, a functor

$$O_a : (A_0\textit{-modules of finite type}) \to (A_0\textit{-modules of finite type}).$$

(2) For each surjection $A' \to A$ with kernel M, an A_0-module of finite type, a class $o_a \in O_a(M)$ which is zero if and only if there exists a lifting of a to A'.

This data is further required to be functorial and linear in (A_0, M). We obtain such a theory by defining $O_a(M) := \mathrm{Ext}^1(La^*L_{\mathcal{Y}/S}, M)$, and taking for each $A' \to A$ as in (2.8.2) the class $o_a \in \mathrm{Ext}^1(La^*L_{\mathcal{Y}/S}, M)$ to be the class obtained from (2.7.1). Note that it follows from the construction of the cotangent complex that the homology groups of $La^*L_{\mathcal{Y}/S}$ are coherent. From this and standard properties of cohomology it follows that the A_0-modules $\mathrm{Ext}^1(La^*L_{\mathcal{Y}/S}, M)$ are of finite type and that the additional conditions ([Ar74], 4.1) on the obstruction theory hold.

A corollary to the above discussion is the following analogy of ([Ko96], Theorem I.2.17):

2.9. **Corollary.** *Let $\mathcal{F} : \mathcal{C}/S \to \mathcal{Y}/S$ be a representable proper morphism from a twisted curve to a smooth Deligne-Mumford stack $\mathcal{Y}/S$. Let F be a field and $s : \mathrm{Spec}\, F \to S$ a morphism. Let $\mathcal{C}^F$ and $\mathcal{Y}^F$ be the fibers over* $\mathrm{Spec}\,(F)$ *and $f : \mathcal{C}^F \to \mathcal{Y}^F$ the morphism induced by $\mathcal{F}$. Assume that S is equidimensional at s. Then the dimension of every irreducible component of $\underline{\mathrm{Hom}}_S(\mathcal{C}, \mathcal{Y})$ at $[f]$ is at least*

$$H^0(\mathcal{C}, f^*(T_{\mathcal{Y}^F})) - H^1(\mathcal{C}, f^*(T_{\mathcal{Y}^F})) + \dim_s S.$$

Furthermore, if $H^1(\mathcal{C}, f^*(T_{\mathcal{Y}^F})) = 0$, then $\underline{\mathrm{Hom}}_S(\mathcal{C}, \mathcal{Y})$ is smooth over S at $[f]$.

Proof. Because of (2.7.2), we know that the tangent space of $\underline{\mathrm{Hom}}_s(\mathcal{C}^F, \mathcal{Y}^F)$ (the fiber of $\underline{\mathrm{Hom}}_S(\mathcal{C}, \mathcal{Y})$ at s) is isomorphic to $H^0(\mathcal{C}, f^*(T_{\mathcal{Y}^F}))$. Let (Q, m_Q) be the local ring of a point of an étale altas of $\underline{\mathrm{Hom}}_S(\mathcal{C}, \mathcal{Y})$, whose image is $[f]$. It is a quotient of a local $\mathcal{O}_{s,S}$-algebra R, which is smooth over S of relative dimension $h^0(\mathcal{C}, f^*(T_{\mathcal{Y}^F}))$. Let K be the kernel. Then $K \subset (m_S R, m_R^2)$ because Q and R have the same relative tangent space.

Let $B = R/m_R K$ and m_B be the corresponding maximal ideal. The kernel J of $B \to Q$ is isomorphic to $K/m_R K$. Because of (2.8), we know J has a largest quotient $J \to J^m$ such that the image of

$$o \in \mathrm{Ext}^1(f^*(\Omega_{\mathcal{Y}^F}), J) \cong H^1(\mathcal{C}, f^*T_{\mathcal{Y}^F}) \otimes J$$

in $H^1(\mathcal{C}, f^*T_{\mathcal{Y}^F}) \otimes J^m$ is 0. So since the obstruction theory is linear, we have

$$\dim J^m \geq \dim J - h^1(\mathcal{C}, f^*T_{\mathcal{Y}^F}).$$

On the other hand, $\dim J = \dim K/m_R K$, and $J^m = 0$ since

$$0 \to I^m \to J \to J^m \to 0,$$

which implies, by the definition of J^m, the split of $B/I^m \to B/J \cong Q$, which then implies $J^m = 0$. So

$$\text{number of generators of } K = \dim K/m_R K \leq h^1(\mathcal{C}, f^*T_{\mathcal{Y}^F})$$

□

3. Smoothing

Henceforce, we focus on the n-pointed twisted stable maps of genus 0, i.e. objects in $\mathcal{K}_{0,n}(\mathcal{M}, d)$. In the classical case, to solve the problems relating to rational curves, one of the main tools is the comb construction and its deformations (cf. [KMM92b], [Ko96] and [GHS03]). Here we generalize this technique to n-pointed twisted stable maps, and also to the case that the handle is reducible.

3.1. **Definition.** *A comb with m teeth* is a twisted nodal curve $\mathcal{T}$ over k with the following properties:

1. $\mathcal{T} = \mathcal{D} \cup_{1 \leq i \leq m} C^i$, where each C^i is isomorphic to $\mathbb{P}^1$;
2. Every C_i intersects $\mathcal{D}$ in a single point x_i in the smooth scheme locus of $\mathcal{D}$ such that $x_i \neq x_j$ for $i \neq j$; and
3. $C^i \cap C^j = \emptyset$.

We call $\mathcal{D}$ the *handle* of $\mathcal{T}$ and C^i the *teeth* of $\mathcal{T}$.

3.2. **Construction.** Let $(f : \mathcal{C} \to \mathcal{M}, s_1, s_2, ..., s_n)$ be an n-pointed genus 0 twisted stable map of degree $d(> 0)$ over k. Now we assume

1. $\mathcal{M}$ is a Deligne-Mumford smooth stack,
2. there is a strongly rationally connected dense open subscheme $M^0 \subset \mathcal{M}$, and
3. if $\mathcal{D}_j$ is an irreducible component of $\mathcal{C}$ such that $\pi(\mathcal{D}_j)$ is not a point in $\mathcal{M}$, then $\pi(\mathcal{D}_j) \cap M^0 \neq \emptyset$.

Let $\mathcal{D}_1$, $\mathcal{D}_2$,..., $\mathcal{D}_s$ be all the components of $\mathcal{C}$, which are not mapped to $\mathcal{M}$ constantly. Because of assumption (3), on each $\mathcal{D}_i$, we can choose m_i general (scheme) points p^i_1, p^i_2,..., $p^i_{m_i}$, which are mapped into M^0.
By assumption (2), we can attach very free curves to $\mathcal{D}$ as teeth. More precisely, we can assume that there exist $m = \sum_{i=1}^s m_i$ 1-pointed very free curves

$$(g^i_j : \mathbb{P}^1 \to M^0, r^i_j),$$

such that

$$g^i_j(r^i_j) = f(p^i_j), \forall i, j.$$

Let $W^i_j \subset p^i_j \times \mathbb{A}^m$ be defined by the equation $y^i_j = 0$ where the coordinates of $\mathbb{A}^m$ are $(y^1_1, ..., y^1_{m_1}, ..., y^i_{m_i})$. Then W^i_j's are disjoint codimension two substacks of $\mathcal{C} \times \mathbb{A}^m$, which are in fact schemes. Let $h : \mathcal{S} \to \mathcal{C} \times \mathbb{A}^m \to \mathbb{A}^m$ be obtained by blowing up all W^i_j. Gluing the morphisms f and g^i_j, we get a morphism $F : \mathcal{S}_0 = h^{-1}(\{0\}) \to \mathcal{M}$. Each $s_i : \Sigma_i (\cong [\operatorname{Spec}(k)/G]) \to \mathcal{C}$ can be lifted to a natural morphism $s'_i : \Sigma_i \times \mathbb{A}^m \to \mathcal{S}$ because $\mathcal{S} \to \mathcal{C} \times \mathbb{A}^m$ induces an isomorphism over the image of $s_i \times \mathbb{A}^m$.

3.3. **Proposition.** *(Notations as above.) There are a morphism* $\gamma : Z \to \mathbb{A}^m$ *(*Z *a curve) with a commutative diagram*

$$\begin{tikzcd}
\mathcal{T} \arrow[r, "\gamma_Z"] \arrow[d, "h_Z"'] & \mathcal{S} \arrow[d, "h"'] \\
Z \arrow[r, "\gamma"] & \mathbb{A}^m
\end{tikzcd}$$

where $\mathcal{T}$ *is a family of subcombs over* Z *containing the image of* $t_i (:= s'_i \times \gamma) : \Sigma_i \times Z \to \mathcal{S} \times_{\mathbb{A}^m} Z$*, and a representable morphism* $\sigma : \mathcal{T} \to \mathcal{M}$ *such that:*

(a) *if we denote the coarse moduli map as* $\pi_{\mathcal{M}} : \mathcal{M} \to M$*, then for each* i*, the image of* $\pi_{\mathcal{M}} \circ \sigma \circ t_i(\Sigma_i \times Z)$ *is a point in* M*;*

(b) *the restriction* γ_η *of* $\gamma_Z : \mathcal{T} \to \mathcal{S}$ *to the fiber over a general point* $\eta \in Z$ *induces an 1-isomorphism between* $h_Z^{-1}(\eta)$ *and* $\mathcal{C}$*;*

(c) *the fiber* $\mathcal{T}_0$ *of* $\mathcal{T}$ *over* $\gamma^{-1}(\{0\})$ *has* m' *teeth with*

$$m' \geq m - h^1(\mathcal{C}, (f|_{\mathcal{C}})^* T_{\mathcal{M}}(-\sum s_i)); \text{ and}$$

(d) $\sigma|_{\mathcal{T}_0}$ *is given by the restriction of* F*.*

Proof. Let us consider the natual morphism between Deligne-Mumford stacks

$$\underline{\mathrm{Hom}}_{\mathbb{A}^n}(\mathcal{S}, \mathcal{M} \times \mathbb{A}^n) \to \prod_i \underline{\mathrm{Hom}}(\Sigma_i, \mathcal{M}).$$

$[F]$ gives a point in $\underline{\mathrm{Hom}}_{\mathbb{A}^n}(\mathcal{S}, \mathcal{M} \times \mathbb{A}^n)$. Let

$$\underline{\mathrm{Hom}}_{\mathbb{A}^n}(\mathcal{S}, \mathcal{M} \times \mathbb{A}^n, G)$$

represent the fiber that contains $[F]$. Let c be the codimension of an irreducible component of the image of a small neighborhood of

$$[F] \in \underline{\mathrm{Hom}}_{\mathbb{A}^n}(\mathcal{S}, \mathcal{M} \times \mathbb{A}^n, G) \text{ in } \mathbb{A}^{\mathrm{m}}.$$

There is an irreducible curve Z

$$r : (q \in Z) \to ([F] \in \underline{\mathrm{Hom}}_{\mathbb{A}^n}(\mathcal{S}, \mathcal{M} \times \mathbb{A}^n, G))$$

such that the image of $r(Z)$ in $\mathbb{A}^m$ is contained in at most c coordinate hyperplanes $(y_i = 0)$. By base change we obtain a flat family $\mathcal{W} \to Z$, whose general fiber is a comb with at most c teeth (and the handle 1-isomorphic to $\mathcal{C}$). From the construction, after shrinking Z and removing the irreducible components corresponding to these teeth, we get a family of subcombs $\mathcal{T}$ with a representable

morphism $\sigma : \mathcal{T} \to \mathcal{M}$ satisfying (a) and (b). It also satisfies (c) if in the statement we change $h^1(\mathcal{C}, (f|_{\mathcal{C}})^* T_{\mathcal{M}}(-\sum s_i))$ to c.
To show (c), combining (2.9) and the argument of [Ko96] (II.7.9), we know that there is a smooth morphism $\mathrm{Spec}\,(R) \to \mathbb{A}^m$ such that (the completion of) $\underline{\mathrm{Hom}}_{\mathbb{A}^n}(\mathcal{S}, \mathcal{M} \times \mathbb{A}^n, G)$ is defined in $\mathrm{Spec}\, R$ by at most $h^1(\mathcal{C}, f^* T_{\mathcal{M}}(-\sum s_i))$ equations. Hence,

$$c \leq h^1(\mathcal{C}, f^* T_{\mathcal{M}}(-\sum s_i)).$$

□

For the reader's convenience, let us recall some simple facts on the cohomology of orbi-curves.

3.4. **Lemma.** *Assume* $\mathrm{char}(k) = 0$ *and* $(\mathcal{K}/k, x_i(r_i))$ *is an smooth proper orbi-curve. Let* $\pi : \mathcal{K} \to K$ *give its coarse moduli and* $\mathcal{L}$ *be a vector bundle on* $\mathcal{K}$ *of rank* r. *Then* $L = \pi_*(\mathcal{L})$ *is a vector bundle on* K *of the same rank, and we have the following exact sequence:*

$$0 \to \pi^*(L) \to \mathcal{L} \to \oplus F_i \to 0.$$

Here F_i *are nontrivial sheaves supported on the gerbs* $x_i(r_i)$. *Furthermore, we have*

$$H^i(\mathcal{K}, \mathcal{L}) = H^i(K, L) \text{ for any } i.$$

Proof. Left to the reader. □

From the above discussion, under the 1-isomorphism induced by γ_η, an irreducible component $\mathcal{D}$ is mapped to a point by f if and only if it is mapped to point by σ. To establish the following proposition, besides the assumptions in (3.2), we need to assume that our family $(f : \mathcal{C} \to \mathcal{M}, s_1, s_2, ..., s_n)$ satisfies:

(4) if $p \neq q$, then the image of $\pi_{\mathcal{M}} \circ s_p$ and $\pi_{\mathcal{M}} \circ s_q$ are different points in M.

3.5. **Proposition.** *(Notations as above) Let* $\mathcal{E}_i = \overline{(\mathcal{C} - \sum_{1 \leq j \leq i} \mathcal{D}_j)} \subset \mathcal{C}$. *Denote* $\mathcal{D}_i \cap \mathcal{E}_i = \Xi_i$ *and assume that* $H^1(\mathcal{D}_i, L \otimes f|^*_{\mathcal{D}_i} T_{\mathcal{M}}(-\sum s_i - \Xi_i) = 0$ *for every line bundle* L *on* $\mathcal{D}_i$, *which is a pull back of some line bundle on the coarse space* D_i *with degree at least* d_i. *Let* η *be a general point on* Z, *if*

$$m_i \geq h^1(\mathcal{C}, f^* T_{\mathcal{M}}(-\sum s_i)) + d_i,$$

then we have

$$H^1(\mathcal{T}_\eta, \sigma_\eta^* T_{\mathcal{M}}(-\sum_i t_i(\eta))) = 0,$$

where $\sigma_\eta = \sigma|_{T_\eta}$.

Proof. Because of (3.3.b) we know $\mathcal{T}_\eta$ is 1-isomorphic to $\mathcal{C}$. In particular, we know that the coarse moduli T_η is a tree of rational curves. Below, we will use the same notations to denote the corresponding substacks of $\mathcal{T}_\eta$ and $\mathcal{C}$. Tensor $f^*T_{\mathcal{M}}(-\sum s_i)$ with the exact sequence (set $\mathcal{E}_0 = \mathcal{T}_\eta \cong \mathcal{C}$)

$$0 \to \mathcal{O}_{\mathcal{D}_i}(-\Xi_i) \to \mathcal{O}_{\mathcal{E}_{i-1}} \to \mathcal{O}_{\mathcal{E}_i} \to 0.$$

Combining the argument of [Ko96] (II.7.10), (3.3.c) and the assumption

$$m_i \geq h^1(\mathcal{C}, f^*T_{\mathcal{M}}(-\sum s_i)) + d_i,$$

we know that for each $\mathcal{D}_i \subset T_\eta$,

$$H^1(\mathcal{D}_i, \sigma_\eta|^*_{\mathcal{D}_i} T_{\mathcal{M}}(-\sum s_i - \Xi_i)) = 0.$$

By induction, to prove

$$H^1(\mathcal{T}_\eta, \sigma^*_\eta T_{\mathcal{M}}(-\sum_i t_i(\eta))) = 0,$$

it suffices to verify that $H^1(\mathcal{E}_s, \sigma^*_\eta T_{\mathcal{M}}(-\sum s_i) = 0$. However, $\mathcal{E}_s$ is exactly the disjoint union of all connected components which are mapped to a point. Hence,

$$H^1(\mathcal{E}_s, f^*T_{\mathcal{M}}(-\sum s_i) = H^1(\mathcal{E}_s, \mathcal{O}_{\mathcal{E}_s}(\sum -s_i))^{\oplus d} = H^1(E_s, \mathcal{O}_{E_s}(-\sum s'_i))^{\oplus d},$$

where d is the dimension of $\mathcal{M}$ and (E_s, s'_i) is the coarse moduli space of $(\mathcal{E}_s, s_i)$. Since each component has coarse moduli spaces of arithmetic genus 0 and been mapped to a point on $\mathcal{M}$, by our assumption (4), the restriction of $\mathcal{O}_{E_s}(\sum s'_i)$ to each component is of degree at most 1. Thus we conclude $H^1(E_s, \mathcal{O}_{E_s}(-\sum s'_i)) = 0$.

□

3.6. **Theorem** (Smoothing). *(Notations as above). We can find an n-pointed twisted stable map (over k) $(g : \mathcal{D} \to \mathcal{M}, t_1, ..., t_n)$ of genus 0 such that*

1. *$\mathcal{D}$ is smooth;*
2. *for each i, the induced morphism $f \circ s_i : \Sigma_i \to \mathcal{M}$ has constant image; and*
3. *$H^1(\mathcal{D}, g^*T_{\mathcal{M}}(-\sum s_i)) = 0$.*

In particular, if we assume Σ_i that are all in the scheme locus of $\mathcal{C}$, then $\mathcal{D} \cong \mathbb{P}^1$.

Proof. Let us first assume the existence of such d_i as in (3.5). Then we know there is an n-pointed twisted stable map $(\sigma_\eta : \mathcal{T}_\eta \to \mathcal{M}, s_1, s_2, ..., s_n)$, such that

$$H^1(\mathcal{T}_\eta, \sigma^*_\eta T_{\mathcal{M}}(-s_\eta(\eta))) = 0.$$

We also know T_η can be written as a degeneration of a family of smooth objects (see the remark after (2.3)). Therefore, since the obstruction space vanishes, from the deformation theory, after an étale base change of the family, σ_η can be extended to an n-pointed twisted stable map of this family with constant restriction map on the n marked gerbes. We let a general fiber of this family be our $\mathcal{D}$. Then the vanishing of $H^1(\mathcal{D}, g^*T_{\mathcal{M}}(-\sum s_i))$ is implied by the upper semicontinuity.

For the last statement, recall that $\mathcal{D}/\cup_i \Sigma_i$ is a scheme, so under the assumption that all σ_i are in the scheme locus, $\mathcal{D}$ is actually a scheme.
Finally, to show the existence of d_i, we apply (3.4) to

$$\sigma_\eta|^*_{\mathcal{D}_i} T_{\mathcal{M}}(-\sum s_i - \Xi_i).$$

Its pushforward to D_i is a vector bundle on $\mathbb{P}^1$, which decomposes as $\oplus_k \mathcal{O}(a^i_k)$ for some integers a^i_k. Then we can choose $d_i = \max_k\{-a^i_k, 0\}$. □

From the following proposition,

3.7. **Proposition.** *(*[Vi87]*) A scheme X of finite type over a field of characteristic 0 is the moduli space of some smooth stack $\mathcal{X}$ if and only if X is geometrically unibranch and its normalization has quotient singularities.*

we also have

3.8. **Proposition.** *Let X be a proper variety with only isolated quotient singularities. Assume its smooth locus X^{sm} is rationally connected and there are at most finite points in X^{sm} which are not in the very free locus X^{vf} of X^{sm}. Then in fact X^{sm} is strongly rationally connected.*

Proof. We start the proof with a simple lemma,

3.9. **Lemma.** *Given any two points $x_1, x_2 \in X$, we can find a pointed curve (B, p) and two morphisms $g_i : B \to X$ such that $g_i(p) = x_i$.*

Proof. There exist two pointed curves (B_i, p_i) with two morphisms $h_i : B_i \to X$ such that $h_i(p_i) = x_i$. Consider a curve $B \subset B_1 \times B_2$ which passes through $p = (p_1, p_2)$. So (B, p) and the compositions $h_i \circ \pi_i$ give the sought-for pointed curves and maps, where π_i mean the projections. □

Now for two arbitrary points $x_1, x_2 \in X^{sm}$, we have a curve (B, p) and two morphisms $g_i : B \to X$, such that $g_i(p) = x_i$ as in the above lemma. Shrinking B, we can assume $g_i(B^0) \subset X^{vf}$ $(i = 1, 2)$, where B^0 means $B - \{p\}$. Therefore, (after taking a possible ramified cover), there is morphism

$$f_t : \mathbb{P}^1 \to X^{vf},$$

which connects $g_1(t)$ and $g_2(t)$, where t is a general point of B^0, and

$$H^1(\mathbb{P}^1, f_t^* T_{X^{vf}}(-g_1(t) - g_2(t))) = 0.$$

Considering

$$\mathrm{Hom}_B(B \times \mathbb{P}^1, B \times X^{vf}, g_1, g_2),$$

we know it is smooth over B at $[f_t]$. So there exists an étale morphism $D^0 \to B^0$ from a connected curve, with a lifting

$$D^0 \to \mathrm{Hom}_B(B \times \mathbb{P}^1, B \times X^{vf}, g_1, g_2).$$

This naturally induces a 1-morphism $D^0 \to \mathcal{K}_{0,2}(\mathcal{X}, d)$, where $\mathcal{X}$ is the smooth stack as in (3.7). Then by the properness of $\mathcal{K}_{0,2}(\mathcal{X}, d)$, (after taking a possible quasi-finite cover again) we can complete our family as a 2-pointed twisted stable map

$$(\pi : \mathcal{C} \to D, f_D : \mathcal{C} \to \mathcal{X}, s_1 : D \to \mathcal{C}, s_2 : D \to \mathcal{C})$$

with a limiting object $(f_q : \mathcal{C}_q \to \mathcal{X}, s_1(q), s_2(q))$ such that $f_q(s_i(q)) = x_i (i = 1, 2)$ for some $q \in D$ lying over p. Notice that s_i are in the scheme locus of $\mathcal{C}$.
We apply (3.6) to $(f_q : \mathcal{C}_q \to \mathcal{X}, s_1(q), s_2(q))$ where $\mathcal{M}$ is $\mathcal{X}$ and $\mathcal{M}^0$ is X^{vf} in (3.2). The assumptions (1), (2) and (4) clearly hold in our case. (3) is also true because of our assumption. Thanks to (3.6), we conclude that there is a morphism $g : \mathbb{P}^1 \to \mathcal{X}$, such that $g(\mathbb{P}^1)$ passes through x_1, x_2 and $g^*(T_{\mathcal{M}})(-s_1(q) - s_2(q))$ is semi-positive. Since for $x \neq s_i(q)$, the tangent map of the universal morphism

$$\mathbb{P}^1 \times \underline{\mathrm{Hom}}(\mathbb{P}^1, \mathcal{X}, s_i(q) \to x_i) \to \mathcal{X}$$

at (x, f) is given by the evaluation morphism

$$T_{x,\mathbb{P}^1} \oplus H^0(\mathbb{P}^1, f^*T_{\mathcal{X}}(-s_1(q) - s_2(q))) \to T_{f(x),\mathcal{X}},$$

the universal morphism is smooth at (x, g) for any $x \neq s_i(q)$, which implies it is equidimensional. So we can choose a general member $[H] \in \underline{\mathrm{Hom}}(\mathbb{P}^1, \mathcal{X}, s_i(q) \to x_i)$ such that $H(\mathbb{P}^1)$ does not pass through the finite stacky points in $\mathcal{X}$. Then its image on X gives a free smooth rational curve which is entirely in the smooth locus of X. □

4. C-NONCONTRACTED MINIMAL MODEL PROGRAM

In this section, S is a projective surface with (at worst) quotient singularities. We denote S^{sm} its smooth locus and S^{vf} the (probably empty) very free locus of S^{sm}. Since surface *klt* singularities precisely mean quotient singularities ([KM98]), for any surface S with *klt* singularities, we can construct a smooth proper Deligne-Mumford stack $\mathcal{S}$ which realizes S as its coarse moduli space. To prove (1.4), we use a specific procedure of minimal model program (MMP) for *klt* surface pair to verify the assumptions of (3.8).

4.1. **Definition.** For a pair (S, Δ) and a fixed irreducible curve C in S, a *C-noncontracted MMP* is a MMP for the pair (S, Δ)

$$f : (S, \Delta) = (S_0, \Delta_0) \to (S_1, \Delta_1) \to \cdots \to (S_n, \Delta_n).,$$

such that in each step, we do not contract the image of $[C]$.

Like the usual MMP, the procedure will terminate and in the last step, we will have a surface pair (S_n, Δ_n) satisfying one of the following:

(i) (S_n, Δ_n) is a log del Pezzo surface with $\rho(S) = 1$, and $0 \neq f_*[C] \in N_1(S_n)$;
(ii) (S_n, Δ_n) admits a Fano contraction to a curve B, such that C is finite over B;
(iii) (S_n, Δ_n) is a minimal model; or
(iv) $\rho(S_n) \geq 2$ and $\overline{\mathrm{NE}}(S)$ has only one $(K_S+\Delta)$-negative extremal ray $R_{\geq 0}[C]$.

In Case(iv), if we contract the extremal ray $R_{\geq 0}[C]$, then we have either a birational contraction $(S_n, \Delta_n) \to (S_{n+1}, \Delta_{n+1})$ with $K_{S_{n+1}} + \Delta_{n+1}$ nef or a Fano contraction to a curve.
The reason for us to consider C-noncontracted MMP is encrypted in the following proposition,

4.2. **Proposition.** *Assume S^{sm} is rationally connected. If we run a C-noncontracted MMP for (S, Δ) and we end in case (i) or (ii), then $C \cap S^{vf} \neq \emptyset$.*

Proof. For each case, we will show that there is a free curve $K \subset S_n^{sm}$ which has nonempty intersection with the image of C. Assuming this for the moment, replacing K by its general deformation, we can think of $S \to S_n$ as an isomorphic over a neighborhood of K. So the preimage of K gives a free curve in S^{sm}. Then it is in S^{vf}, provided the rational connectedness of S^{sm}. Hence we conclude $C \cap S^{vf} \neq \emptyset$.
To prove the claim: in Case(i), because of (1.3), S^{sm} contains a very free curve K. But $\rho(S) = 1$ which implies any two curve have a nonempty intersection. In Case(ii), we can choose K to be a general fiber.

□

4.3. **Lemma.** *If $f : (S, \Delta) \to (S', \Delta')$ is an extremal contraction of surface pairs and (S, Δ) is a log del Pezzo surface, then (S', Δ') is also a log del Pezzo surface.*

Proof. We only need to show that for any irreducible curve C on S', $C \cdot (K_{S'} + \Delta') < 0$. This is true because $f^*(K_{S'} + \Delta') = K_S + \Delta + aE$, where E is the exceptional divisor of f and $a < 0$. □

4.4. **Proposition.** *Under the assumption of (1.4) or (1.5), the complement of S^{vf} in S^{sm} consists of finitely many points.*

Proof. We prove the proposition by contradiction. If there is an irreducible curve in the complement, we take its closure in S, and denote it as C. If we run a C-noncontracted MMP, because of (4.2), we can assume it does not end in case (i) or (ii). However, since the Kodaira dimension $\kappa(S, \Delta) = -\infty$, the only remaining possibility is that (S_n, Δ_n) admits a Fano contraction to a curve, such that C is a fiber. In particular $C^2 = 0$. We claim S_n^{sm} is rationally connected. Assuming this for the moment, then there is a very free rational curve $R \subset S_n^{sm}$ such that $R^2 > 0$. Thus we have $C \cdot R > 0$ because of the Hodge Index Theorem.
Now we verify the claim: if (S, Δ) is a log del Pezzo surface, then it follows from Keel-M^{c}kernan's result and (4.3); if S only contains Du Val singularities, then there is an induced morphism $S_i^{sm} \to S_{i+1}^{sm}$ for each step of MMP, because after any extremal contraction, we will get a smooth point. Hence, S_{i+1}^{sm} is rationally connected if so is S_i^{sm}. □

4.5. **Example** (Gorenstein Surfaces)**.** In general, it is not easy to write down the rational curves in the smooth locus explicitly. However, in the case of (1.5), this can be done in the following way: starting with a surface S, we run a minimal model program.

$$S = S_0 \to S_1 \cdots \to S_n \to Y,$$

it ends with a contraction to $Y = \mathbb{P}^1$ or a point.
If $Y = \mathbb{P}^1$, then because we know S_n^{sm} is rationally connected, the data of multiple fibers $(\mathbb{P}^1, \sum m_i p_i)$ satisfy the equation

$$\sum_i (1 - \frac{1}{m_i}) < 2.$$

So it is easy to explicitly construct the rational curves on S_n as a multiple cover of the base $\mathbb{P}^1$.
If Y is a point, then S_n is a rank 1 log del Pezzo surface with at worst Du Val singularities. In this case, there is a classification in [Fu86]. (In fact, by taking the universal cover of S^{sm} and running a minimal model program for its normal closure, we only need to consider the shorter table of all rank 1 log del Pezzo surfaces with (at worst) Du Val singularities and simply connected smooth loci. Such a table is given in [MZ88]).
So once we understand the rational curves on S_n^{sm} well, by induction, the next thing is to investigate what is going on for a birational contraction $f_i : S_i \to S_{i+1}$. Assuming f_i contracts the irreducible curve D, we know $f_i(D) = p \in S_{i+1}$ is a smooth point, and S_i has at most one singular point along D. This singular point is of type A_n and on its minimal resolution $T \to S_i$, the birational transform of D presents a $(-1)-$curve. Because of our assumption on S_{i+1}, we can write down a rational curve $C' \subset S_{i+1}^{sm}$ which connects a general point and p. Futhermore, we can assume that if we write the (reduced) exceptional locus E of $T \to S_{i+1}$ as a union of curves $E_j (1 \leq j \leq n+1)$ such that E_{n+1} is the birational transform of D and $E_j \cap E_{j+1} \neq \emptyset (1 \leq j \leq n)$, then the birational transform C of C' on T intersects E transversely on E_1 at one point and not on any other $E_j (j \neq 1)$.



figure 1

Let us consider a smooth point $q \in D \subset S_i$ and call its preimage in T to be q_T. Let C_1 be a small deformation of C. We consider a chain of rational curves F_0 with three irreducible components and a morphism $g_1 : F_1 \to T$ such that the restriction of g_1 on the three components gives C, E_1 and C_1 (see figure 2). Computing the normal bundle $\mathcal{N}_{F_1}$ of F_1, we know that $\mathcal{N}|E_1 = \mathcal{O}_{E_1}$. Therefore, we can smooth the reducible curve F to a very free curve $C_2 : \mathbb{P}^1 \to T$ (cf. [HT06]) with

$$C_2 \cdot E_2 = C_2 \cdot E = 1.$$

Now we consider a chain of rational curves F_2 with four irreducible components, and a morphism $g_2 : F_2 \to T$ such that the restriction of g_2 on four components gives C, E_1, E_2 and C_2. Since

$$\mathcal{N}_{F_2}|E_j = \mathcal{O}_{E_j}, 1 \leq j \leq 2,$$

again we can smooth F_2 to get a very free curve $C_3 : \mathbb{P}^1 \to T$.



figure 2

Running this argument n times, we get a very free curve C_{n+1}, such that

$$C_{n+1} \cdot E = C_{n+1} \cdot E_{n+1} = 1.$$

Let us consider a chain of rational curves F' with $n+3$ components, and $g : F' \to T$ such that the restriction of g on each components gives C, E_1,...., E_{n+1} and C_{n+1}. Now

$$\mathcal{N}_{F'}|E_j = \mathcal{O}_{E_j}(1 \leq j \leq n) \text{ and } \mathcal{N}_F|E_{n+1} = \mathcal{O}_{n+1}(1).$$

So we can smooth F' to get a very free curve F which still passes q_T. Furthermore, since

$$F \cdot E_j = 0, 1 \leq j \leq n,$$

the image of F in S_i is in the smooth locus of S_i^{sm}.

Proof of (1.7): Let us consider a resolution $g : Y \to S$. Then Y is a smooth rational surface. We have

$$H^i(Y, \mathcal{O}_Y) = 0, \forall i > 0.$$

The Leray spectral sequence says

$$H^i(S, Rg_*^j(\mathcal{O}_Y)) \Rightarrow H^{i+j}(Y, \mathcal{O}_Y).$$

Because S is normal, we have

$$Rg_*^1(Y, \mathcal{O}_Y) \cong H^2(S, \mathcal{O}_S) \cong H^0(S, \omega_S),$$

However, applying the argument of (cf. [Ko96], IV.3.8), we conclude that the rational connectedness of S^0 implies the vanishing of $H^0(S, \omega_S)$.

5. Nonproperness of the complement of the very free locus

In this section, we will present Jason Starr's proof of the statement that the complement of the very free locus in a smooth rationally connected variety does not contain any proper connected component, which is obviously an improvement of (3.8). The idea is applying the new tool of pseudo-ideal sheaves introduced in [SR09] to the degeneration argument discussed in the above sections.

• **Pseudo-ideal Sheaves.** In this subsection we will give a brief introduction to the concept of pseudo-ideal sheaves. For the proofs of the statements and more background, the reader is encouraged to read the original paper.

Let $f : X \to Y$ be a flat, locally finitely presented, proper morphism of algebraic spaces. For every morphism of algebraic spaces, $g : Y' \to Y$, denote by $f_{Y'} : X_{Y'} \to Y'$ the basechange of f.

5.1. **Definition.** For every morphism $g : Y' \to Y$ of algebraic spaces, a flat family of *pseudo-ideal sheaves* of X/Y over Y' is a pair (F, u) consisting of

(1) a Y'-flat, locally finitely presented, quasi-coherent $\mathcal{O}_{X_{Y'}}$-module F, and
(2) an $\mathcal{O}_{X_{Y'}}$-homomorphism $u : F \to \mathcal{O}_{X_{Y'}}$

such that the following induced morphism is zero,

$$u' : \wedge^2 F \to F, f_1 \wedge f_2 \to u(f_1)f_2 - u(f_2)f_1.$$

Denote by $(g : \mathrm{Hilb}_{X/Y} \to Y, C)$ a universal pair of a morphism of algebraic spaces g and a closed subspace C of $\mathrm{Hilb}_{X/Y} \times_Y X$ which is flat, locally finitely presented, and proper over $\mathrm{Hilb}_{X/Y}$, i.e., an object representing the Hilbert functor. Denote by

$$\begin{tikzcd} 0 \arrow[r] & I \arrow[r] & \mathcal{O}_{\mathrm{Hilb}_{X/Y}} \times_Y X \arrow[r] & \mathcal{O}_C \arrow[r] & 0 \end{tikzcd}$$

the natural exact sequence, where

$$u : I \to \mathcal{O}_{\mathrm{Hilb}_{X/Y}} \times_Y X$$

is the ideal sheaf of C in $\mathrm{Hilb}_{X/Y} \times_Y X$. Then the pair (I, u) is a family of pseudo-ideal sheaves of X/Y over $\mathrm{Hilb}_{X/Y}$.

The significance of pseudo-ideal sheaves has to do with restriction to Cartier divisors. Let D be an effective Cartier divisor in X, considered as a closed subscheme of X, and assume D is flat over Y. Denote by $\mathcal{I}_D$ the pullback

$$\mathcal{I}_D := \mathcal{I} \otimes_{\mathcal{O}_X} \mathcal{O}_D$$

on $\mathrm{Hilb}_{X/Y} \times_Y D$. And denote by

$$u_D : \mathcal{I}_D \to \mathcal{O}_{\mathrm{Hilb}_{X/Y} \times_Y D}$$

the restriction of u, then $(\mathcal{I}_D, u_D)$ is a flat family of pseudo-ideal sheaves of D/Y on $\mathrm{Hilb}_{D/Y}$ ([SR09], Proposition 3.4). Denote by

$$\iota_D : \mathrm{Hilb}_{X/Y} \to \mathrm{Pseudo}_{D/Y}$$

the 1-morphism associated to the flat family $(\mathcal{I}_D, u_D)$ of pseudo-ideal sheaves of D/Y over $Hilb_{X/Y}$. This is the *divisor restriction* map.
In the following we discuss the infinitesimal deformation of the divisor restriction map. Let A' be a local Artin $\mathcal{O}_Y$-algebra with maximal ideal m and residue field κ. And let $0 \to J \to A' \to A \to 0$ be an infinitesimal extension, i.e., mJ is zero. Denote by $X_{A'}$, resp. X_A, X_κ, the fiber product of $X \to Y$ with $\operatorname{Spec} A' \to Y$, resp. $\operatorname{Spec} A \to Y$, $\operatorname{Spec} \kappa \to Y$. Let $(F_{A'}, u_{A'})$ be a pseudo-ideal sheaf of D/Y over $\operatorname{Spec} A'$. Denote by (F_A, u_A), resp. (F_κ, u_κ), the restriction of $(F_{A'}, u_{A'})$ to A, resp. to κ. Let $\mathcal{I}_A$ be the ideal sheaf of a flat family C_A of closed subschemes of X/Y over $\operatorname{Spec} A$. Denote by $\mathcal{I}_\kappa$, resp. C_κ, the restriction of $\mathcal{I}_A$ to κ, resp. of C_A to κ. And assume that ι_D sends I_A to (F_A, u_A).

5.2. **Proposition.** *(*[SR09]*, Propsition 3.6) Let n be a nonnegative integer. Assume that C_κ is a regular immersion of codimension n in X_κ, (since X_κ is an algebraic space, in that definition one must replace the Zariski covering by affine schemes by an etale covering by affine schemes).*

There exists an element ω in

$$J \otimes_\kappa H^1(C_\kappa, \mathcal{O}_X(-D) \cdot \operatorname{Hom}_{\mathcal{O}_{C_\kappa}}(\mathcal{I}_\kappa/\mathcal{I}_\kappa^2, \mathcal{O}_{C_\kappa}))$$

which equals 0 if and only if there exists a flat family $C_{A'}$ of closed subschemes of X/Y over $\operatorname{Spec} A'$ *whose restriction to* $\operatorname{Spec} A$ *equals C_A and whose image under ι_D equals $(F_{A'}, u_{A'})$. When it equals 0, the set of such families $C_{A'}$ is naturally a torsor for the κ-vector space*

$$J \otimes_\kappa H^0(C_\kappa, \mathcal{O}_X(-D) \cdot \operatorname{Hom}_{\mathcal{O}_{C_\kappa}}(\mathcal{I}_\kappa/\mathcal{I}_\kappa^2, \mathcal{O}_{C_\kappa})).$$

In particular, if $h^1(C_\kappa, \mathcal{O}_X(-D) \cdot \operatorname{Hom}_{\mathcal{O}_{C_\kappa}}(\mathcal{I}_\kappa/\mathcal{I}_\kappa^2, \mathcal{O}_{C_\kappa}))$ equals 0, then ι_D is smooth at $[C_\kappa]$.

• **Proof of 1.6.** We take a smooth proper closure X of X^0. Let U be the very locus of X^0. Denote $D = X \setminus X^0$ and $Z = X \setminus U$. We claim

5.3. **Proposition.** *Every connected component of Z intersects D.*

Proof. Let $\mu : \widetilde{X} \to X$ be a log resolution of (X, Z). Denote by $\widetilde{D}$ the union of every connected component of $\mu^{-1}(Z)$ whose image intersects D. Every very free curve in $\widetilde{X} \setminus \widetilde{D}$ maps under μ to a very free curve in $X \setminus D$. Thus the union, $\widetilde{U}$, of all very free curves in $\widetilde{X} \setminus \widetilde{D}$ equals $\mu^{-1}(U)$. So the proposition for the original pair (X, X^0) follows from the proposition for the new pair $(\widetilde{X}, \widetilde{X} \setminus \widetilde{D})$. Thus, without loss of generality, assume that Z is an effective Cartier divisor.
By way of contradiction, assume that there exists a connected component W of Z which does not intersect D. Denote the complement of W in Z by V . Let p be a closed point of U. Let B^0 be an affine curve and let

$$\zeta^0 = (\pi^0 : \mathcal{C}^0 \to B^0; \sigma^0 : B^0 \to \mathcal{C}^0, h^0 : \mathcal{C}^0 \to U)$$

be a family of 1-pointed, smooth, genus 0 maps to U such that $h \circ \sigma$ is a constant morphism with image p and such that $\overline{\text{Image}(h)}$ intersects W. By semistable reduction, after replacing B by a finite cover, we may assume that ζ^0 extends to a family of prestable maps,

$$\zeta = (\pi : \mathcal{C} \to B; \sigma : B \to \mathcal{C}; h : \mathcal{C} \to X),$$

over a curve $B = B^0 \cup \{b\}$ such that $h(\mathcal{C}_b)$ intersects W.
Of course this is probably not a family of embedded curves. But that is easily remedied. Choose a projective embedding of B-schemes,

$$(\pi, i) : \mathcal{C} \to B \times_k \mathbb{P}^r.$$

Now define X^{new} to be $X \times_k \mathbb{P}^r$ and define D^{new} to be $D \times_k \mathbb{P}^r$. The union U^{new} of all very free curves in $X^{new} \setminus D^{new}$ equals $U \times \mathbb{P}^r$. And now the map $(h, i) : \mathcal{C} \to X \times_k \mathbb{P}^r$ is a family of embedded curves. Thus, after replacing the original pair (X, D) by the new pair (X^{new}, D^{new}), we may as well assume that the family of curves is embedded.
Denote by C the embedded curve $h(\mathcal{C}_b)$. Denote by G the union of all components of C which intersect U. Now, of course, we apply the "pseudo-ideal sheaf" results from (5.2). Let N be a nonnegative integer such that C is scheme-theoretically contained in $G \cup N \cdot Z$. Denote by Z_N the effective Cartier divisor $N \cdot Z$ of X. Since X is smooth and since Z_N is an effective Cartier divisor, there is an induced "restriction morphism".

$$\iota_{Z_N} : \text{Hilb}_{X_k} \to \text{Pseudo}_{Z_N/k},$$

where $\text{Pseudo}_{Z_N/k}$ is the algebraic stack parameterizing flat families of pseudo-ideal sheaves on Z_N. Every component of G intersects U. Thus by gluing some number m of very free curves in U to each component of G at general points of the component and with general tangent directions, we may form a new nodal, genus 0 curve C_0 in X from C.

5.4. **Lemma.** *If m is sufficiently large, then the morphism ι_{Z_N} is smooth at the Hilbert point $[C_0]$.*

Proof. Because $C \subset G \cup (N \cdot Z)$, $\mathcal{O}_X(-N \cdot Z) \cdot N_{C/X}$ is supported only on G. Then it is a standard argument (see e.g.[GHS03]) that we can add enough very free curves on general points of each component of G in general tangent directions to get a new curve C_0, satisfying

$$h^1(C_0, \mathcal{O}_X(-N \cdot Z) \cdot N_{C_0/X}) = 0,$$

so we can apply (5.2). □

Since Z_N is the disjoint union of $V_N := N \cdot V$ and $W_N := N \cdot W$, there is a canonical isomorphism

$$\text{Pseudo}_{Z_N/k} \cong \text{Pseudo}_{V_N/k} \times_k \text{Pseudo}_{W_N/k}.$$

Denote by h_V the induced morphism

$$\iota_{V_N} \circ \zeta : B \to \mathrm{Hilb}_{X/k} \to \mathrm{Pseudo}_{V_N/k}.$$

i.e., the family of pseudo-ideal sheaves on V_N corresponding to the restriction of the family of curves $\mathcal{C}$ in X. And denote by h_W the constant morphism $h_W : B \to \mathrm{Pseudo}_{W_N/k}$ whose image equals $\iota_{W_N}([C])$, i.e., the pseudo-ideal sheaf on W_N corresponding to the restriction of C to W_N. This defines a morphism

$$(h_V, h_W) : B \to \mathrm{Pseudo}_{Z_N/k}.$$

And by construction $(h_V, h_W)(b)$ equals $\iota_{Z_N}([C_0])$.
Form the 2-fibered product

$$\begin{tikzcd}
H \arrow[r, "q"] \arrow[d, "p"'] & \mathrm{Hilb}_{X/k} \arrow[d, "\iota_{Z_N}"] \\
B \arrow[r] & \mathrm{Pseudo}_{Z_N/k}
\end{tikzcd}$$

where the bottom map is given by (h_V, h_W). Since $(h_V, h_W)(b)$ equals $\iota_{Z_N}([C_0])$, the pair $(b, [C_0])$ is a point of H. Since ι_{Z_N} is smooth at $[C_0]$, the morphism p is smooth at $(b, [C_0])$ by base change. In particular, p is dominant. Therefore there exists a point $(\eta_B, [C_\eta])$ of H mapping under p to the generic point of B.
By construction, $\iota_{V_N}([C_\eta])$ equals $\iota_{V_N}([\mathcal{C}_{\eta_B}])$, where η_B is the generic point of B. But, by construction, the generic member $\mathcal{C}_{\eta_B}$ of the family $\mathcal{C}$ does not intersect V_N. Thus $\iota_{V_N}([\mathcal{C}_B])$ is the "trivial" pseudo-ideal sheaf on V_N, i.e., the identity map $\mathcal{O}_{V_N} \to \mathcal{O}_{V_N}$. Since $\iota_{V_N}([C_\eta])$ is the trivial pseudo-ideal sheaf, also C_η does not intersect V_N.
On the other hand, $\iota_{W_N}([C_\eta])$ equals $\iota_{W_N}([C_0])$, which is nontrivial. Therefore C_η does intersect W_N. Also, since C_η is a generalization of C_0, and since C_0 intersects the open subset U, also C_η intersects the open subset U. Finally, C_η is a generalization of an (embedded) nodal, genus 0 curve in X. Thus C_η is also a at-worst-nodal, genus 0 curve in X. So every irreducible component of C_η is a smooth genus 0 curve. Since C_η is contained in $X \setminus V = (U \cup W)$, and since C_η is a connected curve which intersects both U and W, there exists at least one component of C_η which intersects both U and W. Since this component is contained in $X \setminus V$, in particular it is contained in X^0. But, by the usual comb-smoothing argument, every rational curve in X^0 which intersects the very free locus U is contained in U, contradicting that the component also intersects W. This contradiction proves that, in fact, there is no connected component W which is disjoint from D. □

6. Examples

6.1. **Example.** (Variance of Rational Connectedness in MMP.) In the above arguments, when we run a C-noncontracted MMP for a projective surface S with quotient singularities,

$$S = S_0 \to S_1 \to \cdots \to S_n,$$

under extra assumptions, we can show in each step, the smooth locus S_i^{sm} is rationally connected, provided S_0^{sm} is rationally connected. However, it is somewhat a surprise to know that this is not true in general.

Let us consider $f : S \to \mathbb{P}^1$ a ruled surface obtained in the following way: we start with a Gorenstein ruled surface with two singular fibers over p_1, p_2 of the type $E_1 + 4E_2 + 3E_3 + 2E_4 + 3E_5 + E_6$ contained in the smooth locus and a multiple two fiber over p_3 with $2A_1$ singularities, then we get S by contracting E_1, E_3, E_4 and E_6 on each fiber.



figure 3

where the numbers in the second column mean the self-intersections of the components. Since $1/2 + 2/3 + 2/3 < 2$, we can find multiple sections of S, which are rational curves and meet the reducible fiber in general points of E_5'. This implies that S^{sm} is rationally connected. However, since $K_S \cdot E_5' = -4/15 < 0$, E_5' is a K_S-negative extremal ray, if we contract the E_5' components in the two reduible fibers, then any multiple section $D \to \mathbb{P}^1$ contained in the smooth locus will factor through the orbi-curve

$$D \to (\mathbb{P}^1, p_1(4), p_2(4), p_3(2)).$$

In particular, D is not a rational curve.

In higher dimension, the rational connectedness of the smooth locus is not preserved even for MMP procedures of terminal varieties. In fact, Prokhorov gives an example of a rationally connected variety Y with terminal singularities which admits a Fano contraction to a normal rational surface S with K_S ample and Y(cf. [HX], 5.13). Then we claim

1. Y^{sm} is not rationally connected, and
2. there exists a MMP procedure

$$X = X_0 \to X_1 \to \cdots \to X_n = Y,$$

such that X is a smooth rationally connected variety.

For (1), since $\pi : Y \to S$ is a Fano contraction, and $\mathrm{Sing}(S)$ is of codimension 2, $\pi^{-1}\mathrm{Sing}(S)$ is of codimension at least 2. Therefore, if Y^{sm} is rationally connected, then we can find a very free curve on Y^{sm}, which avoids the subset $\pi^{-1}\mathrm{Sing}(S)$. Then the images of its small deformations give a family of free rational curves on S^{sm}, which contradicts to the assumption that K_S is ample.

For (2), since any variety with terminal singularities is an output of a MMP procedure which starts from a smooth variety (simply reverse the process of resolution of singularities), hence the claim is obvious.

6.2. **Example.** (Gorenstein del Pezzo surfaces in characteristic 2)

In this subsection, we present an example due to Kollár, which is a del Pezzo surface of degree 2 over $\overline{\mathbb{F}}_2$, whose smooth locus does not contain any free rational curves.

We start with $\mathbb{P}^2_{\mathbb{F}_2}$, it has seven $\mathbb{F}_2$-points. There are also seven lines over $\mathbb{F}_2$, each of which passes through precisely three $\mathbb{F}_2$-points. We blow up all these seven $\mathbb{F}_2$-points. The birational transforms of the lines are (-2)-curves. After contracting all these (-2)-curves, we have a Gorenstein del Pezzo surface S of degree 2, which contains seven A_1-points.

$$\begin{tikzcd}
X \arrow[r] \arrow[d] & \mathbb{P}^2_{\mathbb{F}_2} \\
S &
\end{tikzcd}$$

S admits a degree 2 inseparable morphism to $\mathbb{P}^2$, given by the equation $x_3^2 = f_4(x_0, x_1, x_2)$, where f_4 is a degree 4 polynomial with only nondegenerate critical points. S is singular at (x_0, x_1, x_2, x_3) if and only if (x_0, x_1, x_2) is a critical point of S, i.e., the vanishing points of the section $df \in \Omega_1(4)$. For more detailed explanation, see e.g. ([Ko96] V.6.4),

We want to show that there do not exist free curves in S^{sm}. Let I be the idea sheaf of the seven A_1 singularities on S.

6.3. **Lemma.** *There exists an exact sequence of the form*

$$0 \to g^*(I(1)) \to \Omega^1_S \to g^*(\mathcal{O}(-2)) \to 0.$$

Proof. This is essentially the exact sequence of ([Ko95], 9.4). In fact, thanks to the Kollár's construction in [Ko95], we know that C gives a morphism

$$d_C : \mathcal{O}(-4) \to \Omega^1_{\mathbb{P}^2}$$

and an exact sequence

$$0 \to g^*\mathrm{coker}(\mathcal{O}(-4) \to \Omega^1_{\mathbb{P}^2}) \to \Omega^1_S \to g^*(\mathcal{O}(-2)) \to 0.$$

(For the meaning of this exact, see ([Ko95], 9).) Choosing local coordinates to compute, we know

$$(dx, dy)/(xdx + ydy) \cong (x, y)\frac{dx}{y}.$$

We also have the first Chern class

$$c_1(\mathrm{coker}(\mathcal{O}(-4) \to \Omega^1_{\mathbb{P}^2})) = \mathcal{O}(1),$$

so we conclude that

$$\mathrm{coker}(\mathcal{O}(-4) \to \Omega^1_{\mathbb{P}^2}) \cong I(1),$$

□

Now for any rational curve $f : \mathbb{P}^1 \to S^{sm}$, by pulling back the exact sequence by f^*, we can see it is not a free curve.

6.4. **Question.** *It is interesting to ask whether there exists a smooth Fano variety which is defined over a field of characteristic* $p > 0$ *and does not contain any (very) free curve.*

References

[Ar74] Artin, M.; Versal deformations and algebraic stacks. *Invent. Math.* **27** (1974), 165-189.

[AV02] Abramovich, D.; Vistoli, A.; Compactifying the space of stable maps. *J. Amer. Math. Soc.* **15** (2002), no.**1**, 27–75 (electronic).

[Ca92] Campana, F.; Connexité rationelle des variétés de Fano. *Ann. Sci. École Norm. Sup.* **25** (1992) 539-545.

[DM69] Deligne, P.; Mumford, D.; The irreducibility of the space of curves of given genus. *Inst. Hautes tudes Sci. Publ. Math.* No. **36** (1969), 75–109.

[Fu86] Furushima, M.; Singular del Pezzo surfaces and analytic compactifications of 3-dimensional complex affine space $\mathbb{C}^3$. *Nagoya Math. J.* **104** (1986), 1–28.

[GHS03] Graber, T.; Harris, J.; Starr, J.; Families of rationally connected varieties. *J. Amer. Math. Soc.* **16** (2003), no. **1**, 57–67 (electronic).

[HM07] Hacon, C.; McKernan, J.; Shokurov's Rational Connectedness Conjecture, *Duke Math. J.* **138** (2007) no. **1**, 119-136.

[HT06] Hassett, B.; Tschinkel, Y.; Weak approximation over function fields. *Invent. Math.* **163** (2006), no. **1**, 171–190.

[HT08] Hassett, B.; Tschinkel, Y.; Approximation at Places of Bad Reduction for Rationally Connected Varieties, *Pure and Applied Mathematics Quarterly* **4** (2008) no. **3**, 743-766.

[HT] Hassett, B.; Tschinkel, Y.; Weak approximation for hypersurfaces of low degree http://math.rice.edu/∼hassett/papers.html

[HX] Hogadi, A.; Xu, C.; Degenerations of Rationally Connected Varieties, *Trans. Amer. Math. Soc.* **361** (2009), no. **7**, 3931–3949.

[KM98] Kollár, J.; Mori, S.; *Birational Geometry of Algebraic Varieties*, Cambridge Tract. in Math. **134**, Cambridge University Press, Cambridge, 1998.

[KM99] Keel, S.; M^{c}Kernan, J. Rational curves on quasi-projective surfaces. *Mem. Amer. Math. Soc.* **140** (1999), no. **669**.

[KMM92a] Kollár, J.; Miyaoka, Y.; Mori, S.; Rational Connectedness and Boundness of Fano Manifolds, *J. Diff. Geom.* **36** (1992), no. **3**, 765–779.

[KMM92b] Kollár, J.; Miyaoka,Y.; Mori,S.; Rationally connected varieties. *J. Algebraic Geom.* **1** (1992), no. **3**, 429–448.

[Kn08] Knecht, A,; weak approimation for general degree 2 del Pezzo surfaces. arxiv: 0809.9261.

[Ko95] Kollár, J.; Nonrational hypersurfaces. *J. Amer. Math. Soc.* **8** (1995), no. **1**, 241–249.

[Ko96] Kollár, J.; *Rational curves on algebraic varieties.*, Ergeb. Math. Grenz. **3**. Folge, **32**. Springer-Verlag, Berlin, 1996.

[MZ88] Miyanishi, M.; Zhang, D.-Q.; Gorenstein log del Pezzo surfaces of rank one. *J. Algebra* **118** (1988), no. **1**, 63–84.

[Ol06a] Olsson, M.; Deformation theory of representable morphisms of algebraic stacks. *Math. Z.* **253** (2006), no. **1**, 25–62.

[Ol06b] Olsson, M.; $\underline{\text{Hom}}$-stacks and restriction of scalars. *Duke Math. J.* **134** (2006), no. **1**, 139–164.

[Ol07] Olsson, M.; On (log) twisted curves, *Comp. Math.* **143** (2007), 476-494.

[SR09] Starr, J.; Roth, M.; A local-global principle for weak approximation on varieties over function fields. arXiv:0908.0096

[Vi87] Vistoli, A.; Intersection theory on algebraic stacks and on their moduli spaces. *Invent. Math.* **97** (1989), no. **3**, 613–670.

[Xu] Xu, C.; weak approimation for (some) del Pezzo surfaces. arXiv:0902.1765.

[Zh06] Zhang, Q.; Rational connectedness of log $\mathbb{Q}$-Fano varieties. *J. Reine Angew. Math.* **590** (2006), 131–142.

Chenyang Xu
Department of Mathematics, Institute of Advanced Study
Einstein Drive, Princeton, NJ 08540
Email: chenyang@math.ias.edu
Current Address:
Department of Mathematics, Massachusetts Institute of Technology,
Headquarters Office Building 2, Room 236
77 Massachusetts Avenue Cambridge, MA 02139
email cyxu@mit.edu